

# letters to the editor

Please submit letters for the editor's consideration within three weeks of receipt of *Clinical Medicine*. Letters should ideally be limited to 350 words, and sent by email to: [clinicalmedicine@rcplondon.ac.uk](mailto:clinicalmedicine@rcplondon.ac.uk)

## Conversations with Charles

Editor – I wish to thank you for the kind remarks in your editorial (*Clin Med* December 2010, pp 645–6). Shortly after my last manuscript was submitted I realised that contrary to Charles's usual approach I had not gone back to basic principles in my penultimate article and so missed the obvious solution to the problem of establishing an absolute proxy for body bulk.

If one considers the body as a cylinder then bulk is reflected in circumference, and so is essentially two-dimensional. Hence body mass index (BMI) ( $\text{mass}/\text{height}^2$ ) is a good proxy for relative bulk. To correct for the height of the cylinder one has to scale for height by dividing BMI by actual height/median height of the reference population, say 1.7. This gives an absolute proxy for bulk. This can be simplified to Ponderal index ( $\text{mass}/\text{height}^3$ ) multiplied by median height. This standardised BMI would have the great advantage of maintaining the familiar 25 as ideal. Its main benefit would be to avoid underestimate of malnutrition in thin tall people, though in the bulky, where body composition is more relevant, standardised waist measurement might remain as good if not better.

CK CONNOLLY

## Acute decompensated heart failure secondary to thiamine deficiency: often a missed diagnosis

Editor – We read with great interest Saunders *et al's* excellent educational paper (*Clin Med* December 2010 pp 624–7) on malnutrition. Thiamine deficiency leads to dysfunction of

cardiovascular system, commonly known as wet beriberi. This tends to present as acute decompensated heart failure and also with signs of hyperdynamic circulation.<sup>1,2</sup> Cardiac beriberi is usually missed in clinical practice because of the absence of classically described symptoms, such as pedal oedema/anasarca. It has been reported that these patients have ongoing myocardial damage with troponin rise.<sup>3</sup> Patients in the high risk group of thiamine deficiency are likely to be suffering from chronic alcoholism, social isolation or poor dietary intake, including elderly.<sup>4</sup> As the body storage of thiamine is often small, with a high turnover rate and a half life of 10 to 18 days, high risk patients can enter into thiamine deficiency rapidly. Coexisting hypomagnesaemia, a likely result of chronic alcohol abuse, further aggravates the myocardial damaging effect of thiamine deficiency.<sup>5</sup>

These patients are likely to be diagnosed with idiopathic dilated cardiomyopathy as coronary angiography rules out ischaemic cardiomyopathy. From our experience, erythrocyte transketolase levels are under requested for patients presenting with heart failure.

Treatment of such patients includes usual heart failure management and aggressive replenishment of thiamine. This has been shown to have dramatic response to haemodynamic state in a few hours.<sup>1,2</sup>

We suggest that thiamine deficiency should be excluded in malnourished patient's presenting with acute decompensated heart failure.

ASIF KHAN

*Clinical fellow cardiology*

PANKAJ GARG

*Cardiology specialist registrar*

*Aintree Cardiac Centre*

*University Hospital Aintree, Liverpool*

## References

- 1 Rao SN, Chandak GR. Cardiac beriberi: often a missed diagnosis. *J Trop Pediatr* 2010;56:284–5.
- 2 Naidoo DP, Gathiram V, Sadhabiriss A, Hassen F. Clinical diagnosis of cardiac beriberi. *S Afr Med J* 1990;77:125–7.
- 3 Tran H A. Increased troponin I in 'wet' beriberi. *J Clin Pathol* 2009;59:555.
- 4 Russell RM. Vitamin and trace mineral deficiency and excess. In: Braunwald D, Fauci AS, Kasper DL *et al* (eds), *Harrison's principles of internal medicine*, 15th edn. New York: McGraw-Hill, 2001.
- 5 Martin PR, Singleton CK, Hiller-Sturmhofel S. The role of thiamine deficiency in acute brain disease. *Alcohol Res Health* 2003;27:134–42.

## In response

Wet beriberi is fortunately a rare condition in the western world, even allowing for a degree of underreporting and missed diagnosis. Instead, interest focuses on patients with moderate thiamine deficiency and the potential role in chronic heart failure. The incidence of thiamine deficiency in patients with heart failure is reported to range from 13–98% and is exacerbated by malnutrition and loop-diuretics, which promote thiamine excretion.<sup>1–2</sup> There have been mixed results from trials assessing the role of thiamine replacement but one small trial did show some promise with improvements in left ventricular function.<sup>3</sup>

While it is important to obtain a diagnosis if wet beriberi is clinically suspected, given the relative safety of thiamine replacement, it should not be withheld pending laboratory results. Thiamine testing is relatively expensive, has a long laboratory turn around time and blood analysis in acutely unwell patients may not represent tissue availability.

It is also worth remembering that all characteristic deficiency states are the end result of progressive utilisation of micronutrient stores in an attempt to compensate for the shortfall. Identification of one

micronutrient deficiency in an individual from a vulnerable population group should trigger consideration of other subclinical micronutrient deficiencies that may be less apparent.

JOHN SAUNDERS

*Specialist registrar in clinical nutrition  
and gastroenterology*

TREVOR SMITH

*Consultant in clinical nutrition  
and gastroenterology*

*Institute of Human Nutrition,  
Southampton University Hospitals*

## References

- 1 Kwok T, Falconer-Smith JF, Potter JF, Ives DR. Thiamine status of elderly patients with cardiac failure. *Age Ageing* 1992;21:67–71.
- 2 Zenuk C, Healey J, Donnelly J *et al*. Thiamine deficiency in congestive heart failure patients receiving long term furosemide therapy. *Can J Clin Pharmacol* 2003;10:184–8.
- 3 Shimon I, Almog S, Vered Z *et al*. Improved left ventricular function after thiamine supplementation in patients with congestive heart failure receiving long-term furosemide therapy. *Am J Med* 1995;98:485–90.

# Clinical & scientific letters

Letters not directly related to articles published in *Clinical Medicine* and presenting unpublished original data should be submitted for publication in this section. Clinical and scientific letters should not exceed 500 words and may include one table and up to five references.

## Early intervention can influence an outcome – time to introduce an alert system for all cancer patients?

The National Chemotherapy Advisory Group 2009 report<sup>1</sup> emphasised the need for all hospitals with an emergency department (ED) to establish an acute oncology service (AOS). An AOS brings together expertise from oncology disciplines, emergency medicine, general medicine and general surgery. An effective team is one which carefully coordinates care and communicates well.

We have introduced a cost-neutral patient alert system developed by simply linking the hospital number of all known oncology patients to the patient administration system (PAS). Every 24 hours a report is compiled of all tracked patients who have both attended and been admitted via the ED. This report is sent to all members of the acute oncology team (AOT) daily.

The value of such an alert system is epitomised by a case which presented over one weekend: a 40-year-old man known to the local oncologist, having recently undergone hepatic resection of colorectal liver metastases after resection of his primary, following neoadjuvant chemotherapy. He presented with a short history of headaches. He was discharged with simple analgesia with no plans to further investigate or refer back to his oncology team. This seemed highly appropriate as cerebral metastases are not typical of metastatic bowel cancer.

An alert of his attendance was received on Monday morning and followed up by the AOT by calling the patient at home to enquire further about the symptoms. An

outpatient appointment was made for further investigations. Within four days of presentation to the ED, a magnetic resonance imaging (MRI) head scan was performed as his history was suggestive of an acute neurological event. The MRI revealed a two-cm solitary enhancing lesion within the right cerebellar lobe with local mass effect and effacement of the fourth ventricle. He was started on high-dose dexamethasone and his case referred to the specialist neurosurgical service. He underwent a cerebellar metastectomy, histology of which confirmed a sigmoid metastasis. He recovered within days of surgery and has returned to work.

This case illustrates the need for robust systems to ensure that oncology patients are appropriately identified for any new symptoms which may indicate recurrent disease, particularly out of hours. If his isolated cerebellar metastasis was not detected early, as described above, he may have presented following a catastrophic bleed not amenable to a potentially curative procedure.

ANNA GORRINGE

*Core medical trainee, Oncology Department*

JUDITH KING

*Specialist registrar, Oncology Department*

ELIZABETH WHITEHURST

*IM&T Department*

DIANE MURRAY

*Consultant, Radiology Department*

PAULINE LEONARD

*Consultant, Oncology Department*

*Whittington Hospital*

## Reference

- 1 Department of Health. *Chemotherapy services in England, ensuring quality and safety. A report from the National Chemotherapy Advisory Group*. London: DH, 2009. [www.dh.gov.uk/en/Publicationsandstatistics/Publications/DH\\_104500](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/DH_104500)

## Where there is no resident neurologist: a case for a neurology attachment for acute medicine trainees

Acute neurological problems account for 15% of the general unselected medical take and