

- 2 Horridge A. Assessment and investigation of the child with disordered development. *Arch Dis Child Educ Pract Ed* 2011;96:9–20.
- 3 Shapiro K. Assessment of learning disability and cognitive delay. *BMJ Best Practice*, 2017.
- 4 Mithyantha R, Kneen R, McCann E, Gladstone M. Current evidence-based recommendations on investigating children with global developmental delay. *Arch Dis Child* 2017;102:1071–6.
- 5 Hart AR, Sharma R, Atherton M *et al*. Aetiological investigations in early developmental impairment: are they worth it? *Arch Dis Child* 2017;102:1004–13.
- 6 Department of Health. *Valuing people: a new strategy for learning difficulties for the 21st century*. London, DoH 2001.
- 7 Seal AK. The wisdom of investigating early developmental impairment. *Arch Dis Child* 2017;102:999–1000.
- 8 Johannessen J, Nærland T, Hope S *et al*. Parents' Attitudes toward Clinical Genetic Testing for Autism Spectrum Disorder—Data from a Norwegian Sample. *Int J Mol Sci* 2017;18:1078.

## Response

Thank you for your thoughts on our review of an approach to patients with global developmental delay (GDD) and intellectual disability (ID). We were invited to write this article, pitched for physicians (not specialist paediatricians or clinical geneticists) seeing young and older adults with GDD/ID. A generic diagnostic approach to understanding the myriad of causes of GDD/ID is difficult to find due to the non-discrete spectrum between mild, moderate and severe, across multiple domains and this is often a challenging area for general physicians. Our review aimed to help a physician think about ID/GDD in two contexts: firstly, could an identified inherited susceptibility to developmental delay provide insight into future wider health needs and secondly, how the approach to genetic testing in this context is changing. Addressing the points raised in this letter to the editor, the DH definition of learning disability is not misquoted by us. The reference was provided so that readers could read the entire report for more detail. We concur that ethical issues around consent for testing are important. While some families may not seek a diagnosis for their child's developmental delay, our experience suggests that the majority do. Our experience also suggests that families who have children with unexplained GDD/ID often return to clinical genetics services for a diagnosis when their affected children are adults. This may help to inform risk of recurrence of similar problems

in children of unaffected siblings and sometimes for parents to obtain closure. There is no doubt that improved diagnostic and (through the medical literature) prognostic information is often extremely helpful to patients and families in accessing services and healthcare. While all children with GDD should be able to access special education and social care, in practice limited resources means that children with GDD with a specific underlying diagnosis are more likely to be able to access these services than children with GDD without one. In addition, a molecular diagnosis is often key to accessing research studies and to predict the development of any future medical complications. References 4, 5, and 7 here discuss the evidence-based recommendations on investigating children with GDD. These articles were all published in the same issue of *Archives of Disease in Childhood* in November 2017, whereas our review was submitted for publication in May 2017. Chromosomal microarray has been available to patients in Nottingham initially through a research project from the Wellcome Trust Sanger Institute (0.5 Mb BAC array) from 2006 and on a service basis from the Regional Cytogenetics Laboratory from 2009. We have recommended that MRI brain and MRS should be undertaken in patients with GDD/ID who have microcephaly or macrocephaly or abnormal neurological findings, not as a routine investigation. To conclude, we have attempted to summarise an approach to GDD/ID, highlighting that next-generation sequencing in the form of trio whole exome sequencing or whole genome sequencing will likely result in an increasing diagnostic yield in patients with GDD/ID. ■

PRADEEP C VASUDEVAN  
University Hospitals of Leicester NHS Trust,  
Leicester Royal Infirmary, Leicester, UK

MOHNISH SURI  
Nottingham University Hospitals NHS Trust, Nottingham, UK

## Editorial note

We would like to apologise to the authors of *Diagnosis and management of cerebral venous thrombosis (Clin Med 2018;18:75–9.)* for incorrectly listing their job titles and affiliations in the printed version of *Clinical Medicine*. The correct listing is present on all online records of this article.

## Members of the editorial board

Wing Commander Edward Nicol  
Editor-in-chief

Cono Ariti  
Statistical editor

Dr Michael Almond  
Mr Paul Belcher  
Prof Rodger Charlton  
Dr Tim Chevassut

Prof Tahseen Chowdhury  
Dr Nick Cork  
Dr Albert Edwards  
Dr Kate Evans  
Dr Johanna Feary  
Dr Maggie Hammersley  
Dr Dylan Harris  
Prof Brian Hurwitz  
Dr Nicola Jones  
Dr Vikas Kapil

Dr Alexandra Lake  
Dr Tom Levett  
Dr Nicola Lomax  
Prof Philip MacCarthy  
Prof Yash Mahida  
Dr Chris Marguerie  
Dr Ravik Mascarenhas  
Prof Martin McKee  
Dr Andrew Medford  
Dr Rahul Mukherjee

Dr Mehoor Patel  
Dr Gerrard Phillips  
Dr Roby Rakhit  
Dr Jo Rimmer  
Prof Phil Smith  
Dr Angela Star  
Prof Cameron Swift  
Dr Rhys Thomas  
Dr Duncan Wilson  
Dr Juliet Wright