

tionships. Remember most doctors have no business experience and have no consideration of the importance of a 'limited liability partnership'. No book on senior doctoring would be complete without a section on clinical governance and self-regulation and all the usual aspects are covered including again 'finding out what you can do to reduce the risk of adverse events' – 'you are not working in splendid isolation but as part of a team' we are informed, as well as the perverse dangers of intellectual arrogance.

So, is this text likely to help one succeed as a hospital doctor? Well it is certainly not a 'how to' guide, but for those who are determined to succeed and have palpably considered taking action in such a direction, it will act as a useful guide along the way.

Dr PAUL GRANT
Specialist Registrar in Diabetes and Endocrinology
Eastbourne District General Hospital

letters

TO THE EDITOR

Please submit letters for the Editor's consideration within three weeks of receipt of the Journal. Letters should ideally be limited to 350 words, and sent by email to: Clinicalmedicine@rcplondon.ac.uk

Thrombolysis for ischaemic stroke

In their article dealing with thrombolysis treatment for acute ischaemic stroke, Jenkins *et al* (*Clin Med* June 2008 pp 253–8) omitted mentioning two other important flaws in the National Institute of Neurological Disorders and Stroke (NINDS) trial.¹

Firstly, the relevant part of the trial (part 2), that provided the main impetus for this form of treatment, relied on a global test statistic as the primary measure of outcome, and not on incontrovertible hard end-points. The global statistic was itself a composite function of four inter-related neurological scores (the Barthel Index, the Modified Rankin Scale, the National Institutes of Health Stroke Scale, and the Glasgow Outcome Scale). Clinically and statistically significant differences depending on any such composite of inter-related scores must be inherently suspect.

Secondly, part 2 of the trial also required that patients in the control and active treatment groups not receive aspirin in the first 24 hours. Thus, patients treated with

recombinant tissue plasminogen activator (tPA) were compared to controls in receipt of suboptimal standard therapy. To overcome the risk of serious intracranial bleeding due to combined therapy with tPA and aspirin, the investigators should have administered dummy aspirin to the active treatment group and genuine aspirin to the controls.

CR KUMANA
Professor Emeritus
Queen Mary Hospital, Hong Kong

BERNARD MY CHEUNG
Professor of Clinical Pharmacology and
Therapeutics
University of Birmingham

Reference

- 1 Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *N Engl J Med* 1995; 333:1581–7.

In response

We thank Kumana and Cheung for their comments and accept that there are often inherent weaknesses and limitations in neurological scoring systems, although the degree of potential statistical error in this case is debatable in our view. We believe the core aim of the NINDS study was to demonstrate improvement in disability following thrombolysis and, in this respect, we consider that the end-points chosen by the investigators were reasonable. We agree with Kumana and Cheung that the omission of aspirin was not ideal.

PAUL JENKINS
Professor of Medicine
University of Western Australia, Perth

MARTIN TURNER
Specialist Registrar in Neurology
John Radcliffe Hospital, Oxford

PETER JENKINS
Foundation Year 1
Guys and St Thomas' Hospital, London

Post-infective irritable bowel syndrome and dyspepsia

Spiller, in his interesting review (*Clin Med* August 2008 pp 417–9), tabulated nine outbreaks of gastroenteritis followed by irritable bowel syndrome (IBS). However, just as IBS is merely one component of the functional irritable gut syndrome, so some after enteritis had dyspepsia as well as IBS, a phenomenon not detectable using Rome I questionnaires.

The early two Sheffield and three Nottingham studies were uncontrolled. A general practitioner database found that 12 of 318 patients with documented bacterial enteritis developed new IBS, 11 times uninfected controls.¹ In Beijing, 329 of 450 with Shigella gastroenteritis were followed for one to two years as were 243 unaffected siblings or spouses.² Functional bowel disorders developed in 10% who had dysentery <1 week, 31% for 1–2 weeks, and 35% >2 weeks, significantly higher than in controls. A year after a Shigella outbreak in Seoul, incidence of IBS was 17%, but 6% in age- and sex-matched healthy volunteers.³ Two years after a Canadian *Escherichia coli*/*Campylobacter jejuni* outbreak, IBS incidence was 24% but 10% in controls.⁴

Dyspepsia was investigated in two studies. Three months after the Nottingham *C. jejuni* infections 30 with IBS scored 9 for indigestion and 8 for diarrhoea – significantly higher than 28 'controls' (6 and 3) or 40 healthy volunteers (3 and 3).⁵ A one-year follow-up study of a salmonella outbreak in 1,243 Catalan villagers gave a cumulative risk of IBS of 11.6% v 1.5% in controls; the risk of dyspepsia (pain, discomfort, fullness, nausea) was 17.2% v 3.3% in controls.⁶ At 12 months there were 18 villagers who had developed only dyspepsia, 9 with only IBS and 15 with both.

Spiller reviewed possible mechanisms for low-level residual intestinal inflammation in new IBS after gastroenteritis. However infective gastroenteritis might be followed by another mysterious 'functional' alimentary disorder: dyspepsia.

JEREMY HUGH BARON
Honorary Professorial Lecturer,
Mt Sinai School of Medicine, New York

References

- 1 Rodríguez LAG, Ruigómez A. Increased risk of irritable bowel syndrome after bacterial gastroenteritis: cohort study. *BMJ* 1999;318:565–6.
- 2 Wang L-H, Fang X-C, Pan G-Z. Bacillary dysentery as a causative factor of irritable bowel syndrome and its pathogenesis. *Gut* 2004;53:1096–01.
- 3 Ji S, Park H, Lee D *et al*. Post-infectious irritable bowel syndrome in patients with Shigella infection. *J Gastroenterol Hepatol* 2005;20:381–6.
- 4 Marshall JK, Thabane M, Garg AX, Clark WF, Collins SM. Incidence and epidemiology of irritable bowel syndrome after a large waterborne outbreak of bacterial dysentery. *Gastroenterology* 2006; 131:445–50.
- 5 Dunlop SP, Jenkins D, Neal KR, Spiller RC.

Relative importance of enterochromaffin cell hyperplasia, anxiety and depression in postinfectious IBS. *Gastroenterology* 2003; 125:1651–9.

- 6 Mearin F, Pérez-Oliveras M, Perello O *et al*. Dyspepsia and irritable bowel syndrome after a Salmonella gastroenteritis outbreak: one-year follow-up cohort study. *Gastroenterology* 2005;129:98–104.

In response

Professor Baron correctly draws our attention to the fact that IBS is not unique in the functional gastrointestinal diseases (FGIDs) in being precipitated by an infectious insult. The outbreak of salmonellosis in Spain was followed by functional dyspepsia in nearly 1 in 5 of infected individuals.¹ A large series of patients with functional dyspepsia from a tertiary referral centre reported a presumed post-infectious origin in 17% of the patients. This might be important to recognise for management purposes since they had more early satiety, weight loss, nausea, and vomiting than those without an infectious onset and also more frequently had impaired accommodation.²

As well as aiding management, recognising post-infectious cases of FGIDs are particularly useful to aid understanding. Since they have a defined onset it is easier to work out the direction of causality of the multiple risk factors. Prospective studies of post-infectious IBS clearly show that developing IBS is dependent on psychological central factors as well as peripheral gastrointestinal factors. One would suspect functional dyspepsia would be similar in this respect.

ROBIN SPILLER
Professor of Gastroenterology
Wolfson Digestive Diseases Centre
University Hospital, Nottingham

References

- 1 Mearin F, Perez-Oliveras M, Perello A *et al*. Dyspepsia and irritable bowel syndrome after a Salmonella gastroenteritis outbreak: one-year follow-up cohort study. *Gastroenterology* 2005;129:98–104.
- 2 Tack J, Demedts I, Dehondt G *et al*. Clinical and pathophysiological characteristics of acute-onset functional dyspepsia. *Gastroenterology* 2002;122: 1738–47.

■ CONVERSATIONS WITH CHARLES

Screen for malignant arrhythmias in all young (and elderly) athletes?

I read the other day that Italians are required to have a medical examination, including an electrocardiogram (ECG), if they are to participate in sport organised by a third party. As we finished our relatively gentle game of tennis at the club I mentioned this to Charles.

'So if we were in Italy we would have had our ECG checked as we walked onto the court this evening!'

'That's going a bit far, Charles' *I replied.*

'Of course it is, but it makes my point. A difficult question that applies to all screening procedures is "How often?". I might have an ECG today and have a silent coronary tomorrow.'

'That's true but what you can do is look statistically at the law of diminishing returns from repeated

examinations and set an appropriate interval in the light of this.'

'True, but I thought that exercise benefits those with heart disease. Might not the person be discouraged and the club, or at least its insurance company, be apprehensive if the ECG did suggest coronary abnormality?'

'Any competent physician should be able to advise his patient appropriately and so effectively indemnifying the club against serious mishaps.'

'That only covers those with known disease, Coe! What about those with occult disease? Surely they are better to exercise in happy ignorance?'

'There is certainly an argument that there is likely to be net benefit in encouraging exercise in those with

Clin Med
2008;8:557–8